

# Key RMI DHS Indicators



	Residence			Education		
	RMI	Urban	Rural	No education/ primary	Secondary	More than secondary
<b>Marriage and fertility</b>						
1. Women aged 20–24 married by age 18 (%)	26.3	na	na	na	na	na
2. Men aged 20–24 married by age 18 (%)	11.8	na	na	na	na	na
3. Total fertility rate (children per woman)	4.5	4.1	5.2	4.4	4.7	3.2
4. Women aged 15–19 who were already mothers or pregnant at the time of the survey	26.3	20.1	42.8	30.1	25.1	*
5. Median age at first birth for women aged 25–49	20.2	20.5	19.3	18.6	20.3	22.3
6. Married women (15–49 years) with 2 living children wanting no more children (%)	52.4	54.1	48.6	54.7	51.3	53.9
<b>Family planning (currently married women aged 15–49)</b>						
<b>Current use of family planning</b>						
7. Any method (%)	44.6	42.7	48.3	50.1	43.2	39.9
8. Any modern method (%)	42.4	41.2	44.7	47.2	41.6	36.1
8a. Female sterilization (%)	25.0	25.8	23.2	30.5	23.5	20.5
8b. Male sterilization (%)	0.2	0.0	0.7	0.3	0.2	0.0
8c. Injectables (%)	9.2	6.5	14.6	10.6	9.2	6.5
8d. Pill (%)	2.7	3.1	2.1	0.6	3.1	5.8
8e. Condom (%)	1.4	1.9	0.4	1.7	1.3	1.4
<b>Unmet need for family planning</b>						
9. Total unmet need	8.1	7.3	9.6	9.0	7.6	8.9
9a. For spacing (%)	3.0	3.1	2.8	2.5	2.9	4.6
9b. For limiting (%)	5.1	4.2	6.8	6.4	4.7	4.3
<b>Infant and child mortality</b>						
<b>For the 10-year period preceding the survey</b>						
10. Neonatal mortality	23	21	27	14	27	*
11. Infant mortality rate	33	30	37	28	35	*
12. Under 5 mortality rate	46	44	49	47	47	*
<b>For the 5-year period preceding the survey</b>						
13. Neonatal mortality	15	na	na	na	na	na
14. Infant mortality rate	21	na	na	na	na	na
15. Under 5 mortality rate	37	na	na	na	na	na
<b>Maternal and child health</b>						
<b>Maternity care (for births in the last 3 years)</b>						
16. Mothers who had at least 4 antenatal care visits for their last birth (%)	77.1	76.6	78.1	na	na	na
17. Births assisted by a doctor/nurse/other health personnel	94.1	96.7	89.5	89.3	95.0	99.2
18. Births that took place in a hospital or health facility	85.1	94.7	67.9	75.4	86.7	97.0
19. Mothers who received postnatal care within 2 days of delivery for their last birth	63.6	61.1	68.0	61.2	66.6	49.4
<b>Child immunization and vitamin A supplementation</b>						
20a. Children 12–23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	34.3	43.6	13.4	35.6	34.2	*
20b. Children 12–23 months who have received BCG (%)	69.5	68.2	72.4	67.6	71.0	*
20c. Children 12–23 months who have received 3 doses of polio vaccine (%)	45.9	58.3	17.9	46.4	45.4	*
20d. Children 12–23 months who have received 3 doses of DPT vaccine (%)	47.8	60.6	18.8	44.6	49.1	*
20e. Children 12–23 months who have received measles vaccine (%)	54.1	55.9	50.0	49.5	55.8	*
21. Children 12–35 months who received a vitamin A supplement in the last 6 months (%)	53.3	57.3	46.1	57.0	53.3	44.4
<b>Treatment of childhood diseases (children under 3 years)</b>						
22. Children with diarrhea in the last 2 weeks who received ORS	38.2	38.9	37.0	(51.1)	28.6	*
23. Children with diarrhea in the last 2 weeks taken to a health facility (%)	51.4	40.7	69.4	(63.4)	46.1	*
24. Children with fever in the last 2 weeks taken to a health facility (%)	62.8	55.3	(78.3)	(71.9)	57.6	*
<b>Child feeding practices and nutritional status of children</b>						
25. Children under 3 years breastfed within 1 hour of birth (%)	72.5	69.3	78.4	73.6	71.6	76.3
26. Children 0–5 months exclusively breastfed (%)	27.3	na	na	na	na	na
27. Children 6–8 months receiving solid or semi-solid food and breastmilk (%)	86.5	na	na	na	na	na
28. Children under 3 years who are stunted (%)	na	na	na	na	na	na
29. Children under 3 years who are wasted (%)	na	na	na	na	na	na
30. Children under 3 years who are underweight (%)	na	na	na	na	na	na
<b>Knowledge of HIV and AIDS among ever married adults (aged 15–49)</b>						
31. Women who have heard of AIDS (%)	95.8	97.0	93.4	91.1	97.0	100
32. Men who have heard of AIDS (%)	96.6	96.0	98.0	95.1	96.8	98.8
33. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	73.1	76.5	66.0	69.7	74.3	74.5
34. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	89.7	90.8	87.1	86.6	90.3	92.6
35. Men who had high risk sex in the past 12 months (%)	39.4	41.6	35.0	42.8	40.5	30.1
36. Women who had high risk sex in the past 12 months (%)	18.1	17.2	19.8	15.9	19.4	15.6
<b>Women's empowerment</b>						
37. Currently married women who participate in decision making (household decisions, own health care and visits to relatives)	56.8	57.1	56.2	56.5	56.6	58.2
38. Ever married women who have ever experienced spousal violence (%)	29.6	30.0	28.4	26.5	32.1	24.2

NOTE: Figures in parentheses are based on 25–49 unweighted cases

\* Indicates a figure based on fewer than 25 unweighted cases

na: not available